

-continued

```

<400> SEQUENCE: 263

gacccccctcc accccgcctc                               20

<210> SEQ ID NO 264
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 264

ggtgagtgag tgtgtgcgtg                               20

<210> SEQ ID NO 265
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 265

gtgcggcaag agcttcagcc                               20

<210> SEQ ID NO 266
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 266

gaagtcgtgc tgcttcattgt ggtcggggta gcggctgaag cactgcacgc cataggtcag      60
ggtggtcacg aggggtgggccc agggcacggg cagcttgccc                               100

<210> SEQ ID NO 267
<211> LENGTH: 127
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 267

gccacctacg gcaagctgac cctgaagttc atctgcacca ccggcaagct gcccggtgcc      60
tggccccacc tcgtgaccac cctgacctat ggcgtgcagt gcttcagccg ctaccccgac     120
cacatga                                           127

```

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method of inhibiting off-target cleavage of a DNA molecule by a first guide RNA-endonuclease complex, wherein the first guide RNA-endonuclease complex comprises a first guide RNA comprising a nucleotide target recognition sequence complementary to a first target sequence, the method comprising:

contacting the DNA molecule with a second guide RNA-endonuclease complex, wherein the second guide RNA-endonuclease complex comprises a second guide RNA comprising a nucleotide target recognition sequence with 16 or fewer nucleotides and is complementary to a second target sequence in the DNA molecule, wherein the second target sequence is different from the first target sequence but the second

target sequence is capable of cleavage at a measurable rate by the first guide RNA-endonuclease complex.

2. The method of claim 1, further comprising contacting the DNA molecule with the first guide RNA-endonuclease complex, and wherein second guide RNA-endonuclease complex is contacted to the DNA molecule prior to or simultaneously with the first guide RNA-endonuclease complex.

3. The method of claim 2, wherein the first guide RNA-endonuclease complex and the second guide RNA-endonuclease complex are contacted to the DNA molecule at a ratio of about 20:1 to about 1:20.

4. The method of claim 1, wherein the second target sequence differs from the first target sequence by 0-10 nucleotide mismatches.

5. The method of claim 1, wherein the first guide RNA-endonuclease complex comprises a first endonuclease and